

Список литературы к разделу

Арилуглеводородный рецептор (АНР) и ИММУНИТЕТ

Литература к части I раздела об арильном углеводородном рецепторе АНР

1. Nebert, D.W. Proposed role of drug-metabolizing enzymes: Regulation of steady state levels of the ligands that effect growth, homeostasis, differentiation, and neuroendocrine functions. *Mol. Endocrinol.* 1991, 5, 1203–1214. [[Google Scholar](#)] [[CrossRef](#)]
2. Rannug, A.; Rannug, U. The tryptophan derivative 6-formylindolo[3,2-b]carbazole, FICZ, a dynamic mediator of endogenous aryl hydrocarbon receptor signaling, balances cell growth and differentiation. *Crit. Rev. Toxicol.* 2018, 48, 555–574. [[Google Scholar](#)] [[CrossRef](#)]
3. Rannug, A.; Rannug, U.; Rosenkranz, H.; Winqvist, L.; Westerholm, R.; Agurell, E.; Grafstrom, A.-K. Certain photooxidized derivatives of tryptophan bind with very high affinity to the Ah receptor and are likely to be endogenous signal substances. *J. Biol. Chem.* 1987, 262, 15422–15427. [[Google Scholar](#)] [[PubMed](#)]
4. Rannug, U.; Rannug, A.; Sjöberg, U.; Li, H.; Westerholm, R.; Bergman, J. Structure elucidation of two tryptophan-derived, high affinity Ah receptor ligands. *Chem. Biol.* 1995, 2, 841–845. [[Google Scholar](#)] [[CrossRef](#)]
5. Smirnova, A.; Wincent, E.; Vikström Bergander, L.; Alsberg, T.; Bergman, J.; Rannug, A.; Rannug, U. Evidence for New Light-Independent Pathways for Generation of the Endogenous Aryl Hydrocarbon Receptor Agonist FICZ. *Chem. Res. Toxicol.* 2016, 29, 75–86. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
6. Oberg, M.; Bergander, L.; Håkansson, H.; Rannug, U.; Rannug, A. Identification of the tryptophan photoproduct 6-formylindolo[3,2-b]carbazole, in cell culture medium, as a factor that controls the background aryl hydrocarbon receptor activity. *Toxicol. Sci.* 2005, 85, 935–943. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
7. Diani-Moore, S.; Labitzke, E.; Brown, R.; Garvin, A.; Wong, L.; Rifkind, A.B. Sunlight generates multiple tryptophan photoproducts eliciting high efficacy CYP1A induction in chick hepatocytes and In Vivo. *Toxicol. Sci.* 2006, 90, 96–110. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
8. Fritsche, E.; Schäfer, C.; Calles, C.; Bernsmann, T.; Bernshausen, T.; Wurm, M.; Hubenthal, U.; Cline, J.E.; Hajimiragha, H.; Schroeder, P.; et al. Lightning up the UV response by identification of the arylhydrocarbon receptor as a cytoplasmatic target for ultraviolet B radiation. *Proc. Natl. Acad. Sci. USA* 2007, 104, 8851–8856. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
9. Cibrian, D.; Saiz, M.L.; de la Fuente, H.; Sanchez-Diaz, R.; Moreno-Gonzalo, O.; Jorge, I.; Ferrarini, A.; Vazquez, J.; Punzon, C.; Fresno, M.; et al. CD69 controls the uptake of L-tryptophan through LAT1-CD98 and AhR-dependent secretion of IL-22 in psoriasis. *Nat. Immunol.* 2016, 17, 985–996. [[Google Scholar](#)] [[CrossRef](#)]
10. Magiatis, P.; Pappas, P.; Gaitanis, G.; Mexia, N.; Melliou, E.; Galanou, M.; Blachos, C.; Stathopoulou, K.; Skaltsounis, A.L.; Marselos, M.; et al. Malassezia yeasts produce a collection of exceptionally potent activators of the Ah (dioxin) receptor detected in diseased human skin. *J. Investig. Dermatol.* 2013, 133, 2023–2030. [[Google Scholar](#)] [[CrossRef](#)]
11. Schallreuter, K.U.; Salem, M.A.E.; Gibbons, N.C.J.; Maitland, D.J.; Marsch, E.; Elwary, S.M.A.; Healey, A.R. Blunted epidermal L-tryptophan metabolism in vitiligo affects immune response and ROS scavenging by Fenton chemistry, part 2: Epidermal H₂O₂/ONOO(-)-mediated stress in vitiligo hampers indoleamine 2,3-dioxygenase and aryl hydrocarbon receptor-mediated immune response signaling. *FASEB J.* 2012, 26, 2471–2485. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
12. Zhang, M.; Fan, H.; Tan, S.; Tang, Q.; Liu, X.; Zuo, D.; Liao, Y.; Nan, Z.; Tan, C. The Chinese medicinal herb decoction QRZSLXF enhances anti-inflammatory effect in TNBS-induced colitis via balancing Th17/Tregs differentiation. *J. Ethnopharmacol.* 2020, 251, 112549. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
13. Wincent, E.; Amini, N.; Luecke, S.; Glatt, H.; Bergman, J.; Crescenzi, C.; Rannug, A.; Rannug, U. The suggested physiologic aryl hydrocarbon receptor activator and cytochrome P4501 substrate 6-formylindolo[3,2-b]carbazole is present in humans. *J. Biol. Chem.* 2009, 284, 2690–2696. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
14. Esser, C.; Rannug, A. The aryl hydrocarbon receptor in barrier organ physiology, immunology, and toxicology. *Pharmacol. Rev.* 2015, 67, 259–279. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]

15. Stockinger, B.; di Meglio, P.; Gialitakis, M.; Duarte, J.H. The aryl hydrocarbon receptor: Multitasking in the immune system. *Annu. Rev. Immunol.* 2014, 32, 403–432. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
16. Nobs, S.P.; Tuganbaev, T.; Elinav, E. Microbiome diurnal rhythmicity and its impact on host physiology and disease risk. *EMBO Rep.* 2019, 20, e47129. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
17. Piper, C.J.M.; Rosser, E.C.; Oleinika, K.; Nistala, K.; Krausgruber, T.; Rendeiro, A.F.; Banos, A.; Drozdov, I.; Villa, M.; Thomson, S.; et al. Aryl Hydrocarbon Receptor Contributes to the Transcriptional Program of IL-10-Producing Regulatory B Cells. *Cell Rep.* 2019, 29, 1878–1892.e7. [[Google Scholar](#)] [[CrossRef](#)]
18. Monteleone, I.; Rizzo, A.; Sarra, M.; Sica, G.; Sileri, P.; Biancone, L.; Macdonald, T.T.; Pallone, F.; Monteleone, G. Aryl hydrocarbon receptor-induced signals up-regulate IL-22 production and inhibit inflammation in the gastrointestinal tract. *Gastroenterology* 2011, 141, 237–248. [[Google Scholar](#)] [[CrossRef](#)]
19. Qiu, J.; Heller, J.J.; Guo, X.; Chen, Z.-M.E.; Fish, K.; Fu, Y.-X.; Zhou, L. The aryl hydrocarbon receptor regulates gut immunity through modulation of innate lymphoid cells. *Immunity* 2012, 36, 92–104. [[Google Scholar](#)] [[CrossRef](#)]
20. Lamas, B.; Richard, M.L.; Leducq, V.; Pham, H.-P.; Michel, M.-L.; Da Costa, G.; Bridonneau, C.; Jegou, S.; Hoffmann, T.W.; Natividad, J.M.; et al. CARD9 impacts colitis by altering gut microbiota metabolism of tryptophan into aryl hydrocarbon receptor ligands. *Nat. Med.* 2016, 22, 598–605. [[Google Scholar](#)] [[CrossRef](#)]
21. Schiering, C.; Wincent, E.; Metidji, A.; Iseppon, A.; Li, Y.; Potocnik, A.J.; Omenetti, S.; Henderson, C.J.; Wolf, C.R.; Nebert, D.W.; et al. Feedback control of AHR signalling regulates intestinal immunity. *Nature* 2017, 542, 242–245. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
22. Moura-Alves, P.; Puyskens, A.; Stinn, A.; Klemm, M.; Gühlich-Bornhof, U.; Dorhoi, A.; Furkert, J.; Kreuchwig, A.; Protze, J.; Lozza, L.; et al. Host monitoring of quorum sensing during *Pseudomonas aeruginosa* infection. *Science* 2017, 366. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
23. Manzella, C.R.; Ackerman, M.; Singhal, M.; Ticho, A.L.; Ceh, J.; Alrefai, W.A.; Saksena, S.; Dudeja, P.K.; Gill, R.K. Serotonin Modulates AhR Activation by Interfering with CYP1A1-Mediated Clearance of AhR Ligands. *Cell Physiol. Biochem.* 2020, 54, 126–141. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
24. Obata, Y.; Castano, A.; Boeing, S.; Bon-Frauches, A.C.; Fung, C.; Fallesen, T.; Gomez de Agüero, M.; Yilmaz, B.; Lopes, R.; Huseynova, A.; et al. Neuronal programming by microbiota regulates intestinal physiology. *Nature* 2020, 578, 284–289. [[Google Scholar](#)] [[CrossRef](#)]
25. Fu, Z.D.; Selwyn, F.P.; Cui, J.Y.; Klaassen, C.D. RNA-Seq Profiling of Intestinal Expression of Xenobiotic Processing Genes in Germ-Free Mice. *Drug Metab. Dispos.* 2017, 45, 1225–1238. [[Google Scholar](#)] [[CrossRef](#)]
26. Uno, S.; Dragin, N.; Miller, M.L.; Dalton, T.P.; Gonzalez, F.J.; Nebert, D.W. Basal and inducible CYP1 mRNA quantitation and protein localization throughout the mouse gastrointestinal tract. *Free Radic. Biol. Med.* 2008, 44, 570–583. [[Google Scholar](#)] [[CrossRef](#)]
27. Ikuta, T.; Kobayashi, Y.; Kitazawa, M.; Shiizaki, K.; Itano, N.; Noda, T.; Pettersson, S.; Poellinger, L.; Fujii-Kuriyama, Y.; Taniguchi, S.; et al. ASC-associated inflammation promotes cecal tumorigenesis in aryl hydrocarbon receptor-deficient mice. *Carcinogenesis* 2013, 34, 1620–1627. [[Google Scholar](#)] [[CrossRef](#)]
28. Korecka, A.; Dona, A.; Lahiri, S.; Tett, A.J.; Al-Asmakh, M.; Braniste, V.; D'Arienzo, R.; Abbaspour, A.; Reichardt, N.; Fujii-Kuriyama, Y.; et al. Bidirectional communication between the Aryl hydrocarbon Receptor (AhR) and the microbiome tunes host metabolism. *NPJ Biofilms Microbiomes* 2016, 2, 16014. [[Google Scholar](#)] [[CrossRef](#)]
29. Wang, J.; Wang, P.; Tian, H.; Tian, F.; Zhang, Y.; Zhang, L.; Gao, X.; Wang, X. Aryl hydrocarbon receptor/IL-22/Stat3 signaling pathway is involved in the modulation of intestinal mucosa antimicrobial molecules by commensal microbiota in mice. *Innate Immun.* 2018, 24, 297–306. [[Google Scholar](#)] [[CrossRef](#)]
30. Kitajima, S.; Morimoto, M.; Sagara, E.; Shimizu, C.; Ikeda, Y. Dextran sodium sulfate-induced colitis in germ-free IqI/Jic mice. *Exp. Anim.* 2001, 50, 387–395. [[Google Scholar](#)] [[CrossRef](#)]
31. Maslowski, K.M.; Vieira, A.T.; Ng, A.; Kranich, J.; Sierro, F.; Yu, D.; Schilter, H.C.; Rolph, M.S.; Mackay, F.; Artis, D.; et al. Regulation of inflammatory responses by gut microbiota and chemoattractant receptor GPR43. *Nature* 2009, 461, 1282–1286. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
32. Poland, A.; Glover, E.; Kende, A.S. Stereospecific, high affinity binding of 2,3,7,8-tetrachlorodibenzo-p-dioxin by hepatic cytosol. Evidence that the binding species is receptor for induction of aryl hydrocarbon hydroxylase. *J. Biol. Chem.* 1976, 251, 4936–4946. [[Google Scholar](#)] [[PubMed](#)]
33. Wattenberg, L.W.; Leong, J.L.; Strand, P.J. Benzpyrene hydroxylase activity in the gastrointestinal tract. *Cancer Res.* 1962, 22, 1120–1125. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]

34. Rosenberg, D.W. Dietary modulation of cytochrome P450 in the small intestinal epithelium. *Pharmacology* 1991, 43, 36–46. [[Google Scholar](#)] [[CrossRef](#)]
35. Loub, W.D.; Wattenberg, L.W.; Davis, D.W. Aryl hydrocarbon hydroxylase induction in rat tissues by naturally occurring indoles of cruciferous plants. *J. Natl. Cancer Inst.* 1975, 54, 985–988. [[Google Scholar](#)]
36. Park, J.Y.; Bjeldanes, L.F. Organ-selective induction of cytochrome P-450-dependent activities by indole-3-carbinol-derived products: Influence on covalent binding of benzo(a)pyrene to hepatic and pulmonary DNA in the rat. *Chem. Biol. Interact.* 1992, 83, 235–247. [[Google Scholar](#)] [[CrossRef](#)]
37. Bjeldanes, L.F.; Kim, J.Y.; Grose, K.R.; Bartholomew, J.C.; Bradfield, C.A. Aromatic hydrocarbon responsiveness-receptor agonists generated from indole-3-carbinol In Vitro and In Vivo: Comparisons with 2,3,7,8-tetrachlorodibenzo-p-dioxin. *Proc. Natl. Acad. Sci. USA* 1991, 88, 9543–9547. [[Google Scholar](#)] [[CrossRef](#)]
38. Wincent, E.; Bengtsson, J.; Mohammadi Bardbori, A.; Alsberg, T.; Luecke, S.; Rannug, U.; Rannug, A. Inhibition of cytochrome P450-dependent clearance of the endogenous agonist FICZ as a mechanism for activation of the aryl hydrocarbon receptor. *Proc. Natl. Acad. Sci. USA* 2012, 109, 4479–4484. [[Google Scholar](#)] [[CrossRef](#)]
39. Westerink, W.M.; Stevenson, J.C.; Schoonen, W.G. Pharmacologic profiling of human and rat cytochrome P450 1A1 and 1A2 induction and competition. *Arch. Toxicol.* 2008, 82, 909–921. [[Google Scholar](#)] [[CrossRef](#)]
40. Shimada, T.; Tanaka, K.; Takenaka, S.; Murayama, N.; Martin, M.V.; Foroozesh, M.K.; Yamazaki, H.; Guengerich, F.P.; Komori, M. Structure-function relationships of inhibition of human cytochromes P450 1A1, 1A2, 1B1, 2C9, and 3A4 by 33 flavonoid derivatives. *Chem. Res. Toxicol.* 2010, 23, 1921–1935. [[Google Scholar](#)] [[CrossRef](#)]
41. Blacher, E.; Levy, M.; Tatirovsky, E.; Elinav, E. Microbiome-Modulated Metabolites at the Interface of Host Immunity. *J. Immunol.* 2017, 198, 572–580. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
42. Agus, A.; Planchais, J.; Sokol, H. Gut Microbiota Regulation of Tryptophan Metabolism in Health and Disease. *Cell Host Microbe* 2018, 23, 716–724. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
43. Gutiérrez-Vázquez, C.; Quintana, F.J. Regulation of the Immune Response by the Aryl Hydrocarbon Receptor. *Immunity* 2018, 48, 19–33. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
44. Gao, J.; Xu, K.; Liu, H.; Liu, G.; Bai, M.; Peng, C.; Li, T.; Yin, Y. Impact of the Gut Microbiota on Intestinal Immunity Mediated by Tryptophan Metabolism. *Front. Cell. Infect. Microbiol.* 2018, 8, 13. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
45. Roager, H.M.; Licht, T.R. Microbial tryptophan catabolites in health and disease. *Nat. Commun.* 2018, 9, 3294. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
46. Wikoff, W.R.; Anfora, A.T.; Liu, J.; Schultz, P.G.; Lesley, S.A.; Peters, E.C.; Siuzdak, G. Metabolomics analysis reveals large effects of gut microflora on mammalian blood metabolites. *Proc. Natl. Acad. Sci. USA* 2009, 106, 3698–3703. [[Google Scholar](#)] [[CrossRef](#)]
47. Zheng, X.; Xie, G.; Zhao, A.; Zhao, L.; Yao, C.; Chiu, N.H.; Zhou, Z.; Bao, Y.; Jia, W.; Nicholson, J.K.; et al. The footprints of gut microbial-mammalian co-metabolism. *J. Proteome. Res.* 2011, 10, 5512–5522. [[Google Scholar](#)] [[CrossRef](#)]
48. Zelante, T.; Iannitti, R.G.; Cunha, C.; Luca, A.; Giovannini, G.; Pieraccini, G.; Zecchi, R.; D'Angelo, C.; Massi-Benedetti, C.; Fallarino, F.; et al. Tryptophan catabolites from microbiota engage aryl hydrocarbon receptor and balance mucosal reactivity via interleukin-22. *Immunity* 2013, 39, 372–385. [[Google Scholar](#)] [[CrossRef](#)]
49. Hashimoto, T.; Perlot, T.; Rehman, A.; Trichereau, J.; Ishiguro, H.; Paolino, M.; Sigl, V.; Hanada, T.; Hanada, R.; Lipinski, S.; et al. ACE2 links amino acid malnutrition to microbial ecology and intestinal inflammation. *Nature* 2012, 487, 477–481. [[Google Scholar](#)] [[CrossRef](#)]
50. Vyhliďalová, B.; Krasulová, K.; Pečinková, P.; Marcalíková, A.; Vrzal, R.; Zemánková, L.; Vančo, J.; Trávníček, Z.; Vondráček, J.; Karasová, M.; et al. Gut Microbial Catabolites of Tryptophan Are Ligands and Agonists of the Aryl Hydrocarbon Receptor: A Detailed Characterization. *Int. J. Mol. Sci.* 2020, 21, 2614. [[Google Scholar](#)] [[CrossRef](#)]
51. Fukumoto, S.; Toshimitsu, T.; Matsuoka, S.; Maruyama, A.; Oh-Oka, K.; Takamura, T.; Nakamura, Y.; Ishimaru, K.; Fujii-Kuriyama, Y.; Ikegami, S.; et al. Identification of a probiotic bacteria-derived activator of the aryl hydrocarbon receptor that inhibits colitis. *Immunol. Cell Biol.* 2014, 92, 460–465. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
52. Włodarska, M.; Luo, C.; Kolde, R.; d'Hennezel, E.; Annand, J.W.; Heim, C.E.; Krastel, P.; Schmitt, E.K.; Omar, A.S.; Creasey, E.A.; et al. Indoleacrylic Acid Produced by Commensal *Peptostreptococcus* Species Suppresses Inflammation. *Cell Host Microbe* 2017, 22, 25–37. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
53. Kaur, H.; Bose, C.; Mande, S.S. Tryptophan Metabolism by Gut Microbiome and Gut-Brain-Axis: An in silico Analysis. *Front. Neurosci.* 2019, 13, 1365. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
54. Makki, K.; Deehan, E.C.; Walter, J.; Bäckhed, F. The Impact of Dietary Fiber on Gut Microbiota in Host Health and Disease. *Cell Host Microbe* 2018, 23, 705–715. [[Google Scholar](#)] [[CrossRef](#)]

55. Furusawa, Y.; Obata, Y.; Fukuda, S.; Endo, T.A.; Nakato, G.; Takahashi, D.; Nakanishi, Y.; Uetake, C.; Kato, K.; Kato, T.; et al. Commensal microbe-derived butyrate induces the differentiation of colonic regulatory T cells. *Nature* 2013, 504, 446–450. [[Google Scholar](#)] [[CrossRef](#)]
56. Singh, N.; Gurav, A.; Sivaprakasam, S.; Brady, E.; Padia, R.; Shi, H.; Thangaraju, M.; Prasad, P.D.; Manicassamy, S.; Munn, D.H.; et al. Activation of Gpr109a, receptor for niacin and the commensal metabolite butyrate, suppresses colonic inflammation and carcinogenesis. *Immunity* 2014, 40, 128–139. [[Google Scholar](#)] [[CrossRef](#)]
57. Thangaraju, M.; Cresci, G.A.; Liu, K.; Ananth, S.; Gnanaprakasam, J.P.; Browning, D.D.; Mellinger, J.D.; Smith, S.B.; Digby, G.J.; Lambert, N.A.; et al. GPR109A is a G-protein-coupled receptor for the bacterial fermentation product butyrate and functions as a tumor suppressor in colon. *Cancer Res.* 2009, 69, 2826–2832. [[Google Scholar](#)] [[CrossRef](#)]
58. Koh, A.; De Vadder, F.; Kovatcheva-Datchary, P.; Bäckhed, F. From Dietary Fiber to Host Physiology: Short-Chain Fatty Acids as Key Bacterial Metabolites. *Cell* 2016, 165, 1332–1345. [[Google Scholar](#)] [[CrossRef](#)]
59. Zhang, J.; Watson, A.J.; Probst, M.R.; Minehart, E.; Hankinson, O. Basis for the loss of aryl hydrocarbon receptor gene expression in clones of a mouse hepatoma cell line. *Mol. Pharmacol.* 1996, 50, 1454–1462. [[Google Scholar](#)]
60. Gradin, K.; Toftgård, R.; Poellinger, L.; Berghard, A. Repression of dioxin signal transduction in fibroblasts. Identification Of a putative repressor associated with Arnt. *J. Biol. Chem.* 1999, 274, 13511–13518. [[Google Scholar](#)] [[CrossRef](#)]
61. Garrison, P.M.; Rogers, J.M.; Brackney, W.R.; Denison, M.S. Effects of histone deacetylase inhibitors on the Ah receptor gene promoter. *Arch. Biochem. Biophys.* 2000, 374, 161–171. [[Google Scholar](#)] [[CrossRef](#)]
62. Haarmann-Stemann, T.; Bothe, H.; Kohli, A.; Sydlik, U.; Abel, J.; Fritsche, E. Analysis of the transcriptional regulation and molecular function of the aryl hydrocarbon receptor repressor in human cell lines. *Drug Metab. Dispos.* 2007, 35, 2262–2269. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
63. Jin, U.H.; Cheng, Y.; Park, H.; Davidson, L.A.; Callaway, E.S.; Chapkin, R.S.; Jayaraman, A.; Asante, A.; Allred, C.; Weaver, E.A.; et al. Short Chain Fatty Acids Enhance Aryl Hydrocarbon (Ah) Responsiveness in Mouse Colonocytes and Caco-2 Human Colon Cancer Cells. *Sci. Rep.* 2017, 7, 10163. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
64. Zapletal, O.; Tylichová, Z.; Neča, J.; Kohoutek, J.; Machala, M.; Milcová, A.; Pokorná, M.; Topinka, J.; Moyer, M.P.; Hofmanová, J.; et al. Butyrate alters expression of cytochrome P450 1A1 and metabolism of benzo(a)pyrene via its histone deacetylase activity in colon epithelial cell models. *Arch. Toxicol.* 2017, 91, 2135–2150. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
65. Rosser, E.C.; Piper, C.J.M.; Matei, D.E.; Blair, P.A.; Rendeiro, A.F.; Orford, M.; Alber, D.G.; Krausgruber, T.; Catalan, D.; Klein, N.; et al. Microbiota-Derived Metabolites Suppress Arthritis by Amplifying Aryl-Hydrocarbon Receptor Activation in Regulatory B Cells. *Cell Metab.* 2020, 31, 837–851.e10. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
66. Marinelli, L.; Martin-Gallausiaux, C.; Bourhis, J.M.; Béguet-Crespel, F.; Blottière, H.M.; Lapaque, N. Identification of the novel role of butyrate as AhR ligand in human intestinal epithelial cells. *Sci. Rep.* 2019, 9, 643. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
67. Sonnenberg, G.F.; Fouser, L.A.; Artis, D. Border patrol: Regulation of immunity, inflammation and tissue homeostasis at barrier surfaces by IL-22. *Nat. Immunol.* 2011, 12, 383–390. [[Google Scholar](#)] [[CrossRef](#)]
68. Peterson, L.W.; Artis, D. Intestinal epithelial cells: Regulators of barrier function and immune homeostasis. *Nat. Rev. Immunol.* 2014, 14, 141–153. [[Google Scholar](#)] [[CrossRef](#)]
69. Zeng, B.; Shi, S.; Ashworth, G.; Dong, C.; Liu, J.; Xing, F. ILC3 function as a double-edged sword in inflammatory bowel diseases. *Cell Death Dis.* 2019, 10, 315. [[Google Scholar](#)] [[CrossRef](#)]
70. Satoh-Takayama, N.; Vosshenrich, C.A.; Lesjean-Pottier, S.; Sawa, S.; Lochner, M.; Rattis, F.; Mention, J.J.; Thiam, K.; Cerf-Bensussan, N.; Mandelboim, O.; et al. Microbial flora drives interleukin 22 production in intestinal NKp46+ cells that provide innate mucosal immune defense. *Immunity* 2008, 29, 958–970. [[Google Scholar](#)] [[CrossRef](#)]
71. Klose, C.S.; Artis, D. Innate lymphoid cells as regulators of immunity, inflammation and tissue homeostasis. *Nat. Immunol.* 2016, 17, 765–774. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
72. De Agüero, M.G.; Ganal-Vonarburg, S.C.; Fuhrer, T.; Rupp, S.; Uchimura, Y.; Li, H.; Steinert, A.; Heikenwalder, M.; Hapfelmeier, S.; Sauer, U.; et al. The maternal microbiota drives early postnatal innate immune development. *Science* 2016, 351, 1296–1302. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
73. Godinho-Silva, C.; Domingues, R.G.; Rendas, M.; Raposo, B.; Ribeiro, H.; da Silva, J.A.; Vieira, A.; Costa, R.M.; Barbosa-Morais, N.L.; Carvalho, T.; et al. Light-entrained and brain-tuned circadian circuits regulate ILC3s and gut homeostasis. *Nature* 2019, 574, 254–258. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]

74. Domingues, R.G.; Hepworth, M.R. Immunoregulatory Sensory Circuits in Group 3 Innate Lymphoid Cell (ILC3) Function and Tissue Homeostasis. *Front. Immunol.* 2020, 11, 116. [[Google Scholar](#)] [[CrossRef](#)]
75. Kiss, E.A.; Vonarbourg, C.; Kopfmann, S.; Hobeika, E.; Finke, D.; Esser, C.; Diefenbach, A. Natural aryl hydrocarbon receptor ligands control organogenesis of intestinal lymphoid follicles. *Science* 2011, 334, 1561–1565. [[Google Scholar](#)] [[CrossRef](#)]
76. Lee, J.S.; Cella, M.; Colonna, M. AHR and the Transcriptional Regulation of Type-17/22 ILC. *Front. Immunol.* 2012, 3, 10. [[Google Scholar](#)] [[CrossRef](#)]
77. Qiu, J.; Guo, X.; Chen, Z.M.; He, L.; Sonnenberg, G.F.; Artis, D.; Fu, Y.X.; Zhou, L. Group 3 innate lymphoid cells inhibit T-cell-mediated intestinal inflammation through aryl hydrocarbon receptor signaling and regulation of microflora. *Immunity* 2013, 39, 386–399. [[Google Scholar](#)] [[CrossRef](#)]
78. Ito, S.; Chen, C.; Satoh, J.; Yim, S.; Gonzalez, F.J. Dietary phytochemicals regulate whole-body CYP1A1 expression through an arylhydrocarbon receptor nuclear translocator-dependent system in gut. *J. Clin. Investig.* 2007, 117, 1940–1950. [[Google Scholar](#)] [[CrossRef](#)]
79. Wei, Y.D.; Helleberg, H.; Rannug, U.; Rannug, A. Rapid and transient induction of CYP1A1 gene expression in human cells by the tryptophan photoproduct 6-formylindolo[3,2-b]carbazole. *Chem. Biol. Interact.* 1998, 2, 39–55. [[Google Scholar](#)] [[CrossRef](#)]
80. Shimada, T.; Guengerich, F.P. Inhibition of human cytochrome P450 1A1-, 1A2-, and 1B1-mediated activation of procarcinogens to genotoxic metabolites by polycyclic aromatic hydrocarbons. *Chem. Res. Toxicol.* 2006, 19, 288–294. [[Google Scholar](#)] [[CrossRef](#)]
81. Mohammadi-Bardbori, A.; Bengtsson, J.; Rannug, U.; Rannug, A.; Wincent, E. Quercetin, resveratrol, and curcumin are indirect activators of the aryl hydrocarbon receptor (AHR). *Chem. Res. Toxicol.* 2012, 25, 1878–1884. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
82. Schiering, C.; Vonk, A.; Das, S.; Stockinger, B.; Wincent, E. Cytochrome P4501-inhibiting chemicals amplify aryl hydrocarbon receptor activation and IL-22 production in T helper 17 cells. *Biochem. Pharmacol.* 2018, 151, 47–58. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
83. Luecke, S.; Wincent, E.; Backlund, M.; Rannug, U.; Rannug, A. Cytochrome P450 1A1 gene regulation by UVB involves crosstalk between the aryl hydrocarbon receptor and nuclear factor kappaB. *Chem. Biol. Interact.* 2010, 184, 466–473. [[Google Scholar](#)] [[CrossRef](#)]
84. Mohammadi-Bardbori, A.; Bergander, L.V.; Rannug, U.; Rannug, A. NADPH Oxidase-Dependent Mechanism Explains How Arsenic and Other Oxidants Can Activate Aryl Hydrocarbon Receptor Signaling. *Chem. Res. Toxicol.* 2015, 28, 2278–2286. [[Google Scholar](#)] [[CrossRef](#)]
85. Gerbal-Chaloin, S.; Iankova, I.; Maurel, P.; Daujat-Chavanieu, M. Nuclear receptors in the cross-talk of drug metabolism and inflammation. *Drug Metab. Rev.* 2013, 45, 122–144. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
86. Zheng, Y.; Valdez, P.A.; Danilenko, D.M.; Hu, Y.; Sa, S.M.; Gong, Q.; Abbas, A.R.; Modrusan, Z.; Ghilardi, N.; de Sauvage, F.J.; et al. Interleukin-22 mediates early host defense against attaching and effacing bacterial pathogens. *Nat. Med.* 2008, 14, 282–289. [[Google Scholar](#)] [[CrossRef](#)]
87. Pickert, G.; Neufert, C.; Leppkes, M.; Zheng, Y.; Wittkopf, N.; Warntjen, M.; Lehr, H.A.; Hirth, S.; Weigmann, B.; Wirtz, S.; et al. STAT3 links IL-22 signaling in intestinal epithelial cells to mucosal wound healing. *J. Exp. Med.* 2009, 206, 1465–1472. [[Google Scholar](#)] [[CrossRef](#)]
88. Vaishnava, S.; Behrendt, C.L.; Ismail, A.S.; Eckmann, L.; Hooper, L.V. Paneth cells directly sense gut commensals and maintain homeostasis at the intestinal host-microbial interface. *Proc. Natl. Acad. Sci. USA* 2008, 105, 20858–20863. [[Google Scholar](#)] [[CrossRef](#)]
89. Vaishnava, S.; Yamamoto, M.; Severson, K.M.; Ruhn, K.A.; Yu, X.; Koren, O.; Ley, R.; Wakeland, E.K.; Hooper, L.V. The antibacterial lectin RegIIIgamma promotes the spatial segregation of microbiota and host in the intestine. *Science* 2011, 334, 255–258. [[Google Scholar](#)] [[CrossRef](#)]
90. Lindemans, C.A.; Calafiore, M.; Mertelsmann, A.M.; O'Connor, M.H.; Dudakov, J.A.; Jenq, R.R.; Velardi, E.; Young, L.F.; Smith, O.M.; Lawrence, G.; et al. Interleukin-22 promotes intestinal-stem-cell-mediated epithelial regeneration. *Nature* 2015, 528, 560–564. [[Google Scholar](#)] [[CrossRef](#)]
91. Sugimoto, K.; Ogawa, A.; Mizoguchi, E.; Shimomura, Y.; Andoh, A.; Bhan, A.K.; Blumberg, R.S.; Xavier, R.J.; Mizoguchi, A. IL-22 ameliorates intestinal inflammation in a mouse model of ulcerative colitis. *J. Clin. Investig.* 2008, 118, 534–544. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
92. Killig, M.; Glatzer, T.; Romagnani, C. Recognition strategies of group 3 innate lymphoid cells. *Front. Immunol.* 2014, 5, 142. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
93. Yeste, A.; Mascanfroni, I.D.; Nadeau, M.; Burns, E.J.; Tukupah, A.M.; Santiago, A.; Wu, C.; Patel, B.; Kumar, D.; Quintana, F.J. IL-21 induces IL-22 production in CD4+ T cells. *Nat. Commun.* 2014, 5, 3753. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
94. Veldhoen, M.; Hirota, K.; Westendorp, A.M.; Buer, J.; Dumoutier, L.; Renauld, J.C.; Stockinger, B. The aryl hydrocarbon receptor links TH17-cell-mediated autoimmunity to environmental toxins. *Nature* 2008, 453, 106–109. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]

95. Quintana, F.J.; Basso, A.S.; Iglesias, A.H.; Korn, T.; Farez, M.F.; Bettelli, E.; Caccamo, M.; Oukka, M.; Weiner, H.L. Control of T(reg) and T(H)17 cell differentiation by the aryl hydrocarbon receptor. *Nature* 2008, 453, 65–71. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
96. Trifari, S.; Kaplan, C.D.; Tran, E.H.; Crellin, N.K.; Spits, H. Identification of a human helper T cell population that has abundant production of interleukin 22 and is distinct from T(H)-17, T(H)1 and T(H)2 cells. *Nat. Immunol.* 2009, 10, 864–871. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
97. Ramirez, J.M.; Brembilla, N.C.; Sorg, O.; Chicheportiche, R.; Matthes, T.; Dayer, J.M.; Saurat, J.H.; Roosnek, E.; Chizzolini, C. Activation of the aryl hydrocarbon receptor reveals distinct requirements for IL-22 and IL-17 production by human T helper cells. *Eur. J. Immunol.* 2010, 40, 2450–2459. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
98. Ishihara, Y.; Kado, S.Y.; Hoepfer, C.; Harel, S.; Vogel, C.F.A. Role of NF- κ B RelB in Aryl Hydrocarbon Receptor-Mediated Ligand Specific Effects. *Int. J. Mol. Sci.* 2019, 20, 2652. [[Google Scholar](#)] [[CrossRef](#)]
99. Martin, B.; Hirota, K.; Cua, D.J.; Stockinger, B.; Veldhoen, M. Interleukin-17-producing gammadelta T cells selectively expand in response to pathogen products and environmental signals. *Immunity* 2009, 31, 321–330. [[Google Scholar](#)] [[CrossRef](#)]
100. Kimura, A.; Abe, H.; Tsuruta, S.; Chiba, S.; Fujii-Kuriyama, Y.; Sekiya, T.; Morita, R.; Yoshimura, A. Aryl hydrocarbon receptor protects against bacterial infection by promoting macrophage survival and reactive oxygen species production. *Int. Immunol.* 2014, 26, 209–220. [[Google Scholar](#)] [[CrossRef](#)]
101. Ji, T.; Xu, C.; Sun, L.; Yu, M.; Peng, K.; Qiu, Y.; Xiao, W.; Yang, H. Aryl Hydrocarbon Receptor Activation Down-Regulates IL-7 and Reduces Inflammation in a Mouse Model of DSS-Induced Colitis. *Dig. Dis. Sci.* 2015, 60, 1958–1966. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
102. Monteleone, I.; Zorzi, F.; Marafini, I.; Di Fusco, D.; Dinallo, V.; Caruso, R.; Izzo, R.; Franzè, E.; Colantoni, A.; Pallone, F.; et al. Aryl hydrocarbon receptor-driven signals inhibit collagen synthesis in the gut. *Eur. J. Immunol.* 2016, 46, 1047–1057. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
103. Chen, W.; Pu, A.; Sheng, B.; Zhang, Z.; Li, L.; Liu, Z.; Wang, Q.; Li, X.; Ma, Y.; Yu, M.; et al. Aryl hydrocarbon receptor activation modulates CD8 α . *Biomed Pharmacother* 2017, 87, 127–134. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
104. Ma, Y.; Wang, Q.; Yu, K.; Fan, X.; Xiao, W.; Cai, Y.; Xu, P.; Yu, M.; Yang, H. 6-Formylindolo(3,2-b)carbazole induced aryl hydrocarbon receptor activation prevents intestinal barrier dysfunction through regulation of claudin-2 expression. *Chem. Biol. Interact.* 2018, 288, 83–90. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
105. Yu, M.; Wang, Q.; Ma, Y.; Li, L.; Yu, K.; Zhang, Z.; Chen, G.; Li, X.; Xiao, W.; Xu, P.; et al. Aryl Hydrocarbon Receptor Activation Modulates Intestinal Epithelial Barrier Function by Maintaining Tight Junction Integrity. *Int. J. Biol. Sci.* 2018, 14, 69–77. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
106. Yin, J.; Yang, K.; Zhou, C.; Xu, P.; Xiao, W.; Yang, H. Aryl hydrocarbon receptor activation alleviates dextran sodium sulfate-induced colitis through enhancing the differentiation of goblet cells. *Biochem. Biophys. Res. Commun.* 2019, 514, 180–186. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
107. Lozupone, C.A.; Stombaugh, J.I.; Gordon, J.I.; Jansson, J.K.; Knight, R. Diversity, stability and resilience of the human gut microbiota. *Nature* 2012, 489, 220–230. [[Google Scholar](#)] [[CrossRef](#)]
108. Mowat, A.M.; Agace, W.W. Regional specialization within the intestinal immune system. *Nat. Rev. Immunol.* 2014, 14, 667–685. [[Google Scholar](#)] [[CrossRef](#)]
109. Thaiss, C.A.; Levy, M.; Korem, T.; Dohnalová, L.; Shapiro, H.; Jaitin, D.A.; David, E.; Winter, D.R.; Gury-BenAri, M.; Tatirovsky, E.; et al. Microbiota Diurnal Rhythmicity Programs Host Transcriptome Oscillations. *Cell* 2016, 167, 1495–1510.e12. [[Google Scholar](#)] [[CrossRef](#)]
110. Zenewicz, L.A.; Yin, X.; Wang, G.; Elinav, E.; Hao, L.; Zhao, L.; Flavell, R.A. IL-22 deficiency alters colonic microbiota to be transmissible and colitogenic. *J. Immunol.* 2013, 190, 5306–5312. [[Google Scholar](#)] [[CrossRef](#)]
111. Mizoguchi, A.; Yano, A.; Himuro, H.; Ezaki, Y.; Sadanaga, T.; Mizoguchi, E. Clinical importance of IL-22 cascade in IBD. *J. Gastroenterol.* 2018, 53, 465–474. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
112. Brawner, K.M.; Yeramilli, V.A.; Duck, L.W.; Van Der Pol, W.; Smythies, L.E.; Morrow, C.D.; Elson, C.O.; Martin, C.A. Depletion of dietary aryl hydrocarbon receptor ligands alters microbiota composition and function. *Sci. Rep.* 2019, 9, 14724. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
113. Schanz, O.; Chijiwa, R.; Cengiz, S.C.; Majlesain, Y.; Weighardt, H.; Takeyama, H.; Förster, I. Dietary AhR Ligands Regulate AhRR Expression in Intestinal Immune Cells and Intestinal Microbiota Composition. *Int. J. Mol. Sci.* 2020, 21, 3189. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
114. Li, Y.; Innocentin, S.; Withers, D.R.; Roberts, N.A.; Gallagher, A.R.; Grigorieva, E.F.; Wilhelm, C.; Veldhoen, M. Exogenous stimuli maintain intraepithelial lymphocytes via aryl hydrocarbon receptor activation. *Cell* 2011, 147, 629–640. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]

115. Boussenna, A.; Goncalves-Mendes, N.; Joubert-Zakeyh, J.; Pereira, B.; Fraisse, D.; Vasson, M.P.; Texier, O.; Felgines, C. Impact of basal diet on dextran sodium sulphate (DSS)-induced colitis in rats. *Eur. J. Nutr.* 2015, 54, 1217–1227. [[Google Scholar](#)] [[CrossRef](#)]
116. Cui, H.; Cai, Y.; Wang, L.; Jia, B.; Li, J.; Zhao, S.; C hu, X.; Lin, J.; Zhang, X.; Bian, Y.; et al. Berberine Regulates Treg/Th17 Balance to Treat Ulcerative Colitis Through Modulating the Gut Microbiota in the Colon. *Front. Pharmacol.* 2018, 9, 571. [[Google Scholar](#)] [[CrossRef](#)]
117. Ohno, M.; Nishida, A.; Sugitani, Y.; Nishino, K.; Inatomi, O.; Sugimoto, M.; Kawahara, M.; Andoh, A. Nanoparticle curcumin ameliorates experimental colitis via modulation of gut microbiota and induction of regulatory T cells. *PLoS ONE* 2017, 12, e0185999. [[Google Scholar](#)] [[CrossRef](#)]
118. Xuan, H.; Ou, A.; Hao, S.; Shi, J.; Jin, X. Galangin Protects against Symptoms of Dextran Sodium Sulfate-induced Acute Colitis by Activating Autophagy and Modulating the Gut Microbiota. *Nutrients* 2020, 12, 347. [[Google Scholar](#)] [[CrossRef](#)]
119. Alrafas, H.R.; Busbee, P.B.; Nagarkatti, M.; Nagarkatti, P.S. Resveratrol modulates the gut microbiota to prevent murine colitis development through induction of Tregs and suppression of Th17 cells. *J. Leukoc. Biol.* 2019, 106, 467–480. [[Google Scholar](#)] [[CrossRef](#)]
120. Power, K.A.; Lu, J.T.; Monk, J.M.; Lepp, D.; Wu, W.; Zhang, C.; Liu, R.; Tsao, R.; Robinson, L.E.; Wood, G.A.; et al. Purified rutin and rutin-rich asparagus attenuates disease severity and tissue damage following dextran sodium sulfate-induced colitis. *Mol. Nutr. Food Res.* 2016, 60, 2396–2412. [[Google Scholar](#)] [[CrossRef](#)]
121. Martin, D.A.; Bolling, B.W. A review of the efficacy of dietary polyphenols in experimental models of inflammatory bowel diseases. *Food Funct.* 2015, 6, 1773–1786. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
122. Salaritabar, A.; Darvishi, B.; Hadjiakhoondi, F.; Manayi, A.; Sureda, A.; Nabavi, S.F.; Fitzpatrick, L.R.; Nabavi, S.M.; Bishayee, A. Therapeutic potential of flavonoids in inflammatory bowel disease: A comprehensive review. *World J. Gastroenterol.* 2017, 23, 5097–5114. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
123. Busbee, P.B.; Menzel, L.; Alrafas, H.R.; Dopkins, N.; Becker, W.; Miranda, K.; Tang, C.; Chatterjee, S.; Singh, U.; Nagarkatti, M.; et al. Indole-3-; carbinol prevents colitis and associated microbial dysbiosis in an IL-22-dependent manner. *JCI Insight.* 2020, 5. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
124. Goto, H.; Takemura, N.; Ogasawara, T.; Sasajima, N.; Watanabe, J.; Ito, H.; Morita, T.; Sonoyama, K. Effects of fructo-oligosaccharide on DSS-induced colitis differ in mice fed nonpurified and purified diets. *J. Nutr.* 2010, 140, 2121–2127. [[Google Scholar](#)] [[CrossRef](#)]
125. Atarashi, K.; Tanoue, T.; Shima, T.; Imaoka, A.; Kuwahara, T.; Momose, Y.; Cheng, G.; Yamasaki, S.; Saito, T.; Ohba, Y.; et al. Induction of colonic regulatory T cells by indigenous *Clostridium* species. *Science* 2011, 331, 337–341. [[Google Scholar](#)] [[CrossRef](#)]
126. Kaiser, M.M.M.; Pelgrom, L.R.; van der Ham, A.J.; Yazdanbakhsh, M.; Everts, B. Butyrate Conditions Human Dendritic Cells to Prime Type 1 Regulatory T Cells. *Front Immunol.* 2017, 8, 1429. [[Google Scholar](#)] [[CrossRef](#)]
127. Segain, J.P.; Raingeard de la Blétière, D.; Bourreille, A.; Leray, V.; Gervois, N.; Rosales, C.; Ferrier, L.; Bonnet, C.; Blottière, H.M.; Galmiche, J.P. Butyrate inhibits inflammatory responses through NFκB inhibition: Implications for Crohn's disease. *Gut* 2000, 47, 397–403. [[Google Scholar](#)] [[CrossRef](#)]
128. Arpaia, N.; Campbell, C.; Fan, X.; Dikiy, S.; van der Veeken, J.; deRoos, P.; Liu, H.; Cross, J.R.; Pfeffer, K.; Coffey, P.J.; et al. Metabolites produced by commensal bacteria promote peripheral regulatory T-cell generation. *Nature* 2013, 504, 451–455. [[Google Scholar](#)] [[CrossRef](#)]
129. Zhou, L.; Zhang, M.; Wang, Y.; Dorfman, R.G.; Liu, H.; Yu, T.; Chen, X.; Tang, D.; Xu, L.; Yin, Y.; et al. Faecalibacterium prausnitzii Produces Butyrate to Maintain Th17/Treg Balance and to Ameliorate Colorectal Colitis by Inhibiting Histone Deacetylase 1. *Inflamm. Bowel. Dis.* 2018, 24, 1926–1940. [[Google Scholar](#)] [[CrossRef](#)]
130. Thaiss, C.A.; Zeevi, D.; Levy, M.; Zilberman-Schapira, G.; Suez, J.; Tengeler, A.C.; Abramson, L.; Katz, M.N.; Korem, T.; Zmora, N.; et al. Transkingdom control of microbiota diurnal oscillations promotes metabolic homeostasis. *Cell* 2014, 159, 514–529. [[Google Scholar](#)] [[CrossRef](#)]
131. Leone, V.; Gibbons, S.M.; Martinez, K.; Hutchison, A.L.; Huang, E.Y.; Cham, C.M.; Pierre, J.F.; Heneghan, A.F.; Nadimpalli, A.; Hubert, N.; et al. Effects of diurnal variation of gut microbes and high-fat feeding on host circadian clock function and metabolism. *Cell Host Microbe* 2015, 17, 681–689. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
132. Tredger, J.M.; Chhabra, R.S. Circadian variations in microsomal drug-metabolizing enzyme activities in rat and rabbit tissues. *Xenobiotica* 1977, 7, 481–489. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
133. Huang, P.; Ceccatelli, S.; Rannug, A. A study on diurnal mRNA expression of CYP1A1, AHR, ARNT, and PER2 in rat pituitary and liver. *Environ. Toxicol. Pharmacol.* 2002, 11, 119–126. [[Google Scholar](#)] [[CrossRef](#)]

134. Richardson, V.M.; Santostefano, M.J.; Birnbaum, L.S. Daily cycle of bHLH-PAS proteins, Ah receptor and Arnt, in multiple tissues of female Sprague-Dawley rats. *Biochem. Biophys. Res. Commun.* 1998, 252, 225–231. [[Google Scholar](#)] [[CrossRef](#)]
135. Zenewicz, L.A.; Yancopoulos, G.D.; Valenzuela, D.M.; Murphy, A.J.; Stevens, S.; Flavell, R.A. Innate and adaptive interleukin-22 protects mice from inflammatory bowel disease. *Immunity* 2008, 29, 947–957. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]
136. Geremia, A.; Arancibia-Cárcamo, C.V.; Fleming, M.P.; Rust, N.; Singh, B.; Mortensen, N.J.; Travis, S.P.; Powrie, F. IL-23-responsive innate lymphoid cells are increased in inflammatory bowel disease. *J. Exp. Med.* 2011, 208, 1127–1133. [[Google Scholar](#)] [[CrossRef](#)]
137. Frank, D.N.; Amand, A.L.S.; Feldman, R.A.; Boedeker, E.C.; Harpaz, N.; Pace, N.R. Molecular-phylogenetic characterization of microbial community imbalances in human inflammatory bowel diseases. *Proc. Natl. Acad. Sci. USA* 2007, 104, 13780–13785. [[Google Scholar](#)] [[CrossRef](#)]
138. Packey, C.D.; Sartor, R.B. Commensal bacteria, traditional and opportunistic pathogens, dysbiosis and bacterial killing in inflammatory bowel diseases. *Curr. Opin. Infect. Dis.* 2009, 22, 292–301. [[Google Scholar](#)] [[CrossRef](#)]
139. Machiels, K.; Joossens, M.; Sabino, J.; De Preter, V.; Arijis, I.; Eeckhaut, V.; Ballet, V.; Claes, K.; Van Immerseel, F.; Verbeke, K.; et al. A decrease of the butyrate-producing species *Roseburia hominis* and *Faecalibacterium prausnitzii* defines dysbiosis in patients with ulcerative colitis. *Gut* 2014, 63, 1275–1283. [[Google Scholar](#)] [[CrossRef](#)]
140. Qin, J.; Li, Y.; Cai, Z.; Li, S.; Zhu, J.; Zhang, F.; Liang, S.; Zhang, W.; Guan, Y.; Shen, D.; et al. A metagenome-wide association study of gut microbiota in type 2 diabetes. *Nature* 2012, 490, 55–60. [[Google Scholar](#)] [[CrossRef](#)]
141. Consolandi, C.; Turrioni, S.; Emmi, G.; Severgnini, M.; Fiori, J.; Peano, C.; Biagi, E.; Grassi, A.; Rampelli, S.; Silvestri, E.; et al. Behçet's syndrome patients exhibit specific microbiome signature. *Autoimmun. Rev.* 2015, 14, 269–276. [[Google Scholar](#)] [[CrossRef](#)] [[PubMed](#)]

Литература к [части II](#) раздела об арилуглеводородном рецепторе AHR

1. Platzer B, Richter S, Kneidinger D, Waltenberger D, Woisetschlager M, Strobl H. Aryl hydrocarbon receptor activation inhibits in vitro differentiation of human monocytes and Langerhans dendritic cells. *J Immunol.* 2009;183:66–74.
2. Kiss EA, Vonarbourg C, Kopfmann S, Hobeika E, Finke D, Esser C, et al. Natural aryl hydrocarbon receptor ligands control organogenesis of intestinal lymphoid follicles. *Science.* 2011;334:1561–5.
3. Mosmann TR, Cherwinski H, Bond MW, Giedlin MA, Coffman RL. Two types of murine helper T cell clone. I. Definition according to profiles of lymphokine activities and secreted proteins. *J Immunol.* 1986;136:2348–57.
4. Zhu J, Yamane H, Paul WE. Differentiation of effector CD4 T cell populations. *Annu Rev Immunol.* 2010;28:445–89.
5. Rutz S, Ouyang W. Regulation of interleukin-10 and interleukin-22 expression in T helper cells. *Curr Opin Immunol.* 2011;23:605–12.
6. Marshall NB, Vorachek WR, Stepan LB, Mourich DV, Kerkvliet NI. Functional characterization and gene expression analysis of CD4+ CD25+ regulatory T cells generated in mice treated with 2,3,7,8-tetrachlorodibenzo-p-dioxin. *J Immunol.* 2008;181:2382–91.
7. Quintana FJ, Basso AS, Iglesias AH, Korn T, Farez MF, Bettelli E, et al. Control of T(reg) and T(H)17 cell differentiation by the aryl hydrocarbon receptor. *Nature.* 2008;453:65–71.
8. Gandhi R, Kumar D, Burns EJ, Nadeau M, Dake B, Laroni A, et al. Activation of the aryl hydrocarbon receptor induces human type 1 regulatory T cell-like and Foxp3(+) regulatory T cells. *Nat Immunol.* 2010;11:846–53.
9. Apetoh L, Quintana FJ, Pot C, Joller N, Xiao S, Kumar D, et al. The aryl hydrocarbon receptor interacts with c-Maf to promote the differentiation of type 1 regulatory T cells induced by IL-27. *Nat Immunol.* 2010;11:854–61.
10. Veldhoen M, Hirota K, Westendorf AM, Buer J, Dumoutier L, Renauld JC, et al. The aryl hydrocarbon receptor links TH17-cell-mediated autoimmunity to environmental toxins. *Nature.* 2008;453:106–9.
11. Fontenot JD, Gavin MA, Rudensky AY. Foxp3 programs the development and function of CD4+CD25+ regulatory T cells. *Nat Immunol.* 2003;4:330–6.
12. Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. *Science.* 2003;299:1057–61.
13. Brunkow ME, Jeffery EW, Hjerrild KA, Paepfer B, Clark LB, Yasayko SA, et al. Disruption of a new forkhead/winged-helix protein, scurf, results in the fatal lymphoproliferative disorder of the scurfy mouse. *Nat Genet.* 2001;27:68–73.

14. Wildin RS, Ramsdell F, Peake J, Faravelli F, Casanova JL, Buist N, et al. X-linked neonatal diabetes mellitus, enteropathy and endocrinopathy syndrome is the human equivalent of mouse scurfy. *Nat Genet.* 2001;27:18–20.
15. Horwitz DA, Zheng SG, Gray JD. Natural and TGF-beta-induced Foxp3(+)CD4(+) CD25(+) regulatory T cells are not mirror images of each other. *Trends Immunol.* 2008;29:429–35.
16. Weiner HL. Induction and mechanism of action of transforming growth factor-beta-secreting Th3 regulatory cells. *Immunol Rev.* 2001;182:207–14.
17. Roncarolo MG, Gregori S, Battaglia M, Bacchetta R, Fleischhauer K, Levings MK. Interleukin-10-secreting type 1 regulatory T cells in rodents and humans. *Immunol Rev.* 2006;212:28–50.
18. Monteleone I, Rizzo A, Sarra M, Sica G, Sileri P, Biancone L, et al. Aryl hydrocarbon receptor-induced signals up-regulate IL-22 production and inhibit inflammation in the gastrointestinal tract. *Gastroenterology.* 2011;141:237–48, 248 e231.
19. Zhang L, Ma J, Takeuchi M, Usui Y, Hattori T, Okunuki Y, et al. Suppression of experimental autoimmune uveoretinitis by inducing differentiation of regulatory T cells via activation of aryl hydrocarbon receptor. *Invest Ophthalmol Vis Sci.* 2010;51:2109–17.
20. Marshall NB, Kerkvliet NI. Dioxin and immune regulation: emerging role of aryl hydrocarbon receptor in the generation of regulatory T cells. *Ann N Y Acad Sci.* 1183:25–37.
21. Wang L, Zhou GB, Liu P, Song JH, Liang Y, Yan XJ, et al. Dissection of mechanisms of Chinese medicinal formula Realgar-Indigo naturalis as an effective treatment for promyelocytic leukemia. *Proc Natl Acad Sci U S A.* 2008;105:4826–31.
22. Adachi J, Mori Y, Matsui S, Takigami H, Fujino J, Kitagawa H, et al. Indirubin and indigo are potent aryl hydrocarbon receptor ligands present in human urine. *J Biol Chem.* 2001;276:31475–8.
23. Qiu J, Heller JJ, Guo X, Chen ZM, Fish K, Fu YX, et al. The aryl hydrocarbon receptor regulates gut immunity through modulation of innate lymphoid cells. *Immunity.* 2012;36:92–104.
24. Lee JS, Cella M, McDonald KG, Garlanda C, Kennedy GD, Nukaya M, et al. AHR drives the development of gut ILC22 cells and postnatal lymphoid tissues via pathways dependent on and independent of Notch. *Nat Immunol.* 2011;13:144–51.
25. Li Y, Innocenti S, Withers DR, Roberts NA, Gallagher AR, Grigorieva EF, et al. Exogenous stimuli maintain intraepithelial lymphocytes via aryl hydrocarbon receptor activation. *Cell.* 2011;147:629–40.
26. Quintana FJ, Murugaiyan G, Farez MF, Mitsdoerffer M, Tukpah AM, Burns EJ, et al. An endogenous aryl hydrocarbon receptor ligand acts on dendritic cells and T cells to suppress experimental autoimmune encephalomyelitis. *Proc Natl Acad Sci U S A.* 2011;107:20768–73.
27. Funatake CJ, Marshall NB, Stepan LB, Mourich DV, Kerkvliet NI. Cutting edge: activation of the aryl hydrocarbon receptor by 2,3,7,8-tetrachlorodibenzo-p-dioxin generates a population of CD4+CD25+ cells with characteristics of regulatory T cells. *J Immunol.* 2005;175:4184–8.
28. Knutson JC, Poland A. Response of murine epidermis to 2,3,7,8-tetrachlorodibenzo-p-dioxin: interaction of the ah and hr loci. *Cell.* 1982;30:225–34.
29. Chen W, Jin W, Hardegen N, Lei KJ, Li L, Marinos N, et al. Conversion of peripheral CD4+CD25-naive T cells to CD4+CD25+ regulatory T cells by TGF-beta induction of transcription factor Foxp3. *J Exp Med.* 2003;198:1875–86.
30. Kerkvliet NI, Stepan LB, Vorachek W, Oda S, Farrer D, Wong CP, et al. Activation of aryl hydrocarbon receptor by TCDD prevents diabetes in NOD mice and increases Foxp3+ T cells in pancreatic lymph nodes. *Immunotherapy.* 2009;1:539–47.
31. Singh NP, Singh UP, Singh B, Price RL, Nagarkatti M, Nagarkatti PS. Activation of aryl hydrocarbon receptor (AhR) leads to reciprocal epigenetic regulation of FoxP3 and IL-17 expression and amelioration of experimental colitis. *PLoS One.* 2011;6:e23522.
32. Taylor SR, Alexander DR, Cooper JC, Higgins CF, Elliott JI. Regulatory T cells are resistant to apoptosis via TCR but not P2X7. *J Immunol.* 2007;178:3474–82.
33. Stockinger B, Hirota K, Duarte J, Veldhoen M. External influences on the immune system via activation of the aryl hydrocarbon receptor. *Semin Immunol.* 2011;23:99–105.
34. Knutson JC, Poland A. 2,3,7,8-Tetrachlorodibenzo-p-dioxin: failure to demonstrate toxicity in twenty-three cultured cell types. *Toxicol Appl Pharmacol.* 1980;54:377–83.
35. Kimura A, Naka T, Nohara K, Fujii-Kuriyama Y, Kishimoto T. Aryl hydrocarbon receptor regulates Stat1 activation and participates in the development of Th17 cells. *Proc Natl Acad Sci U S A.* 2008;105:9721–6.
36. Benson JM, Shepherd DM. Dietary ligands of the aryl hydrocarbon receptor induce anti-inflammatory and immunoregulatory effects on murine dendritic cells. *Toxicol Sci.* 2011;124:327–38.
37. Nguyen NT, Kimura A, Nakahama T, Chinen I, Masuda K, Nohara K, et al. Aryl hydrocarbon receptor negatively regulates dendritic cell immunogenicity via a kynurenine-dependent mechanism. *Proc Natl Acad Sci U S A.* 2010;107:19961–6.

38. Puccetti P, Grohmann U. IDO and regulatory T cells: a role for reverse signalling and non-canonical NF- κ B activation. *Nature reviews*. 2007;7:817–23.
39. Mellor AL, Munn DH. IDO expression by dendritic cells: tolerance and tryptophan catabolism. *Nature reviews*. 2004;4:762–74.
40. Vogel CF, Goth SR, Dong B, Pessah IN, Matsumura F. Aryl hydrocarbon receptor signaling mediates expression of indoleamine 2,3-dioxygenase. *Biochem Biophys Res Commun*. 2008;375:331–5.
41. Mezrich JD, Fechner JH, Zhang X, Johnson BP, Burlingham WJ, Bradfield CA. An interaction between kynurenine and the aryl hydrocarbon receptor can generate regulatory T cells. *J Immunol*. 2010;185:3190–8.
42. Borsellino G, Kleinewietfeld M, Di Mitri D, Sternjak A, Diamantini A, Giometto R, et al. Expression of ectonucleotidase CD39 by Foxp3⁺ Treg cells: hydrolysis of extracellular ATP and immune suppression. *Blood*. 2007;110:1225–32.
43. Friedman DJ, Kunzli BM, Yi AR, Sevigny J, Berberat PO, Enyoji K, et al. From the Cover: CD39 deletion exacerbates experimental murine colitis and human polymorphisms increase susceptibility to inflammatory bowel disease. *Proc Natl Acad Sci U S A*. 2009;106:16788–93.
44. Bacchetta R, Bigler M, Touraine JL, Parkman R, Tovo PA, Abrams J, et al. High levels of interleukin 10 production in vivo are associated with tolerance in SCID patients transplanted with HLA mismatched hematopoietic stem cells. *J Exp Med*. 1994;179:493–502.
45. Astier AL, Meiffren G, Freeman S, Hafler DA. Alterations in CD46-mediated Tr1 regulatory T cells in patients with multiple sclerosis. *J Clin Invest*. 2006;116:3252–7.
46. Martinez-Forero I, Garcia-Munoz R, Martinez-Pasamar S, Inoges S, Lopez-Diaz de Cerio A, Palacios R, et al. IL-10 suppressor activity and ex vivo Tr1 cell function are impaired in multiple sclerosis. *Eur J Immunol*. 2008;38:576–86.
47. Awasthi A, Carrier Y, Peron JP, Bettelli E, Kamanaka M, Flavell RA, et al. A dominant function for interleukin 27 in generating interleukin 10-producing anti-inflammatory T cells. *Nat Immunol*. 2007;8:1380–9.
48. Pot C, Jin H, Awasthi A, Liu SM, Lai CY, Madan R, et al. Cutting edge: IL-27 induces the transcription factor c-Maf, cytokine IL-21, and the costimulatory receptor ICOS that coordinately act together to promote differentiation of IL-10-producing Tr1 cells. *J Immunol*. 2009;183:797–801.
49. Pot C, Apetoh L, Awasthi A, Kuchroo VK. Molecular pathways in the induction of interleukin-27-driven regulatory type 1 cells. *J Interferon Cytokine Res*. 2010;30:381–8.
50. Pot C, Apetoh L, Kuchroo VK. Type 1 regulatory T cells (Tr1) in autoimmunity. *Semin Immunol*. 2011;23:202–8.
51. Sonnenberg GF, Fouser LA, Artis D. Border patrol: regulation of immunity, inflammation and tissue homeostasis at barrier surfaces by IL-22. *Nat Immunol*. 2011;12:383–90.
52. Grossman WJ, Verbsky JW, Barchet W, Colonna M, Atkinson JP, Ley TJ. Human T regulatory cells can use the perforin pathway to cause autologous target cell death. *Immunity*. 2004;21:589–601.
53. Kemper C, Chan AC, Green JM, Brett KA, Murphy KM, Atkinson JP. Activation of human CD4⁺ cells with CD3 and CD46 induces a T-regulatory cell 1 phenotype. *Nature*. 2003;421:388–92.
54. Magnani CF, Alberigo G, Bacchetta R, Serafini G, Andreani M, Roncarolo MG, et al. Killing of myeloid APCs via HLA class I, CD2 and CD226 defines a novel mechanism of suppression by human Tr1 cells. *Eur J Immunol*. 2011;41:1652–62.
55. Brembilla NC, Ramirez JM, Chicheportiche R, Sorg O, Saurat JH, Chizzolini C. In vivo dioxin favors interleukin-22 production by human CD4⁺ T cells in an aryl hydrocarbon receptor (AhR)-dependent manner. *PLoS One*. 2011;6:e18741.
56. Alam MS, Maekawa Y, Kitamura A, Tanigaki K, Yoshimoto T, Kishihara K, et al. Notch signaling drives IL-22 secretion in CD4⁺ T cells by stimulating the aryl hydrocarbon receptor. *Proc Natl Acad Sci U S A*. 2011;107:5943–8.
57. Liang SC, Tan XY, Luxenberg DP, Karim R, Dunussi-Joannopoulos K, Collins M, et al. Interleukin (IL)-22 and IL-17 are coexpressed by Th17 cells and cooperatively enhance expression of antimicrobial peptides. *J Exp Med*. 2006;203:2271–9.
58. Ghoreschi K, Laurence A, Yang XP, Tato CM, McGeachy MJ, Konkel JE, et al. Generation of pathogenic T(H)17 cells in the absence of TGF- β signalling. *Nature*. 2010;467:967–71.
59. Xu J, Yang Y, Qiu G, Lal G, Wu Z, Levy DE, et al. c-Maf regulates IL-10 expression during Th17 polarization. *J Immunol*. 2009;182:6226–36.
60. Rutz S, Noubade R, Eidenschenk C, Ota N, Zeng W, Zheng Y, et al. Transcription factor c-Maf mediates the TGF- β -dependent suppression of IL-22 production in T(H)17 cells. *Nat Immunol*. 2011;12:1238–45.
61. Opitz CA, Litztenburger UM, Sahm F, Ott M, Tritschler I, Trump S, et al. An endogenous tumour-promoting ligand of the human aryl hydrocarbon receptor. *Nature*. 2011;478:197–203.
62. Zou W. Regulatory T cells, tumour immunity and immunotherapy. *Nature reviews*. 2006;6:295–307.